

## 6. References

- Allan, K. (2006) *Biological effects of secondary salinisation on freshwater macroinvertebrates in Tasmania: the acute salinity toxicity testing of seven macroinvertebrates*. Thesis for Masters of Appl Sci. James Cook University, Townsville, Queensland. 60p.
- ANZECC (1992) *Australian Water Quality Guidelines for Fresh and Marine Waters*. Australian and New Zealand Environment and Conservation Council, Canberra.
- ANZECC (2000) *Australian Water Quality Guidelines for Fresh and Marine Waters*. Australian and New Zealand Environment and Conservation Council, Canberra.
- Barker ,P. (2005). Little Swanport River: vegetation associated with three sites of different sediment and flow characteristics. In Warfe D.M. (2005). *A Framework for Determining Holistic Environmental Water Requirements in Tasmania*. Draft Technical Report. Department of Primary Industries, Water and Environment: Hobart, Tasmania.
- Barmuta, L.A., Chessman, B.C. and Hart, B.T. (1998) *Interpreting the outputs from AUSRIVAS*. Occasional Paper Series No. 02/98. (LWRRDC: Canberra).
- Belbin, L. (1994) PATN Pattern Analysis Package. CSIRO Division of Wildlife and Rangelands Research, Canberra.
- Bendix, J. and Hupp C.R. (2000). Hydrological and geomorphological impacts on riparian plant communities. *Hydrological Processes* **14**: 2977-2990.
- Bessel-Browne, T. (2000). Environmental Health of streams in the Yarra River catchment. EPA Publication 682, Environmental Protection Agency, Melbourne.
- Boulton, A.J., Lake, P.S. (1992) The ecology of two intermittent streams in Victoria, Australia. III. Temporal changes in faunal composition *Freshwater Biology* **27** (1), pp. 123-138.
- Boulton, A.J. and Brock, M.A. (1999) *Australian freshwater ecology: processes and management*. pp 184, Gleneagles Publishing, South Australia.

Buckney, R.T. and Tyler, P.A. (1973) Chemistry of Tasmanian Inland Waters. *Internationale Revue der Gesamten Hydrobiologie*, Volume 58, Number 1, pp67-78.

Bunn, S., Mosisch, T. and Davies, P.M. (1999) Temperature and Light. In Lovett, S. and Price, P. (eds.), *Riparian Land Management Technical Guidelines. Volume One: Principles of Sound Management*. Land and Water Resources Research and Development Corporation, Canberra, ACT.

Chessman, B.C. (1995) Rapid assessment of rivers using macroinvertebrates: a procedure based on habitat -specific sampling, family level identification and a biotic index. *Australian Journal of Ecology*, Vol. 33, pp.122-129.

Close, M.E. and Davies-Colley, R.J. (1990) Baseflow water chemistry in New Zealand rivers 1. Characterisation. *New Zealand Journal of Marine and Freshwater Research*, **24**, pp319-341.

Coysh, J.L., Nichols, S.J., Simpson, J.C., Norris, R.H., Barmuta, L.A., Chessman, B.C. and Blackman, P. (2000). AUSTRALIAN RIVER Assessment System (AUSRIVAS) Web Page

Cummins K.W., (1993) Riparian stream linkages: Instream Issues, in *Ecology and Management of Riparian zones in Australia*. (Eds.S.E. Bunn, B.J. Pusey and P. Price). LWRRDC Occasional Paper Series No. 05/93

Daley, E.A. (2003) *Tasmania's Native Riparian Vegetation*. PhD Thesis, School of Geography and Environmental Studies, University of Tasmania. Hobart, Tasmania.

Davies, P.E. (1989) Relationships between habitat characteristics and population abundance for brown trout, *Salmo trutta* L., and blackfish, *Gadopsis marmoratus* (Rich), in Tasmanian streams. *Australian Journal of Marine and Freshwater Research*, **40**, 341-359.

Davies, P.E. (1994), National River and Management program Monitoring River Health Initiative: River Bio-assessment Manual. Commonwealth Environmental Protection Agency, Canberra.

Davies, J. (1988) *Land systems of Tasmania, Region 6: South, East and Midlands-A Resource Classification Survey*. Department of Agriculture, Tasmania.

Davies, P., Cook, B., Rutherford, K. and Walsh, T. (2004) *Managing high in-stream temperatures using riparian vegetation*. Final report to Land and Water Australia, May 2004, Centre of Excellence in Natural Resource Management, University of Western Australia.

Davies, P.E., Warfe D.M., Parslow J. and Telfer D. (2002). *Environmental Flows for the Lower Derwent River: Final Report to DPIWE*. Freshwater Systems, CSIRO Division of Fisheries and Oceanography, and GECO Stream & Catchment Assessments: Hobart, Tasmania

Davies-Colley, R.J. and Smith, D.G. (2001) Turbidity, Suspended Sediment, And Water Clarity: A Review. *Journal of the American Water Resources Association*, **37**, No 5, pp1085-1101.

DPIWE (1999) *Quality of the Brid River*. Water Assessment Branch, Department of Primary Industries, Water and Environment, Hobart, Tasmania. Technical Report No. WRA 99/09

DPIWE (2003c) *State of the River Report for the Coal River Catchment*. Water Assessment and Planning Branch, Department of Primary Industries, Water and Environment, Hobart. Technical Report No. WAP 03/11

DPIWE (2003) *State of the River Report for the Jordan River Catchment*. Water Assessment and Planning Branch, Department of Primary Industries, Water and Environment, Hobart. Technical Report No. WAP 03/10

DPIWE (2003c) *State of the River Report for the Duck River Catchment*. Water Assessment and Planning Branch, Department of Primary Industries, Water and Environment, Hobart. Technical Report No. WAP 03/08

DPIWE (2003d) *State of Rivers Report for the North Esk Catchment*. Water Assessment and Planning Branch, Department of Primary Industries, Water and Environment, Hobart. Technical Report No. WAP 03/06

DPIWE (2005) *A Framework for Determining Holistic Environmental Water Requirements in Tasmania (DRAFT)*. Water Assessment and Planning Branch, Department of Primary Industries, Water and Environment, Hobart. Technical Report No. WAP 05/02

Dell, M. (2000) *Hydrogeological Setting of Areas Subject to Soil Salinity in Tasmania*. Tasmanian Geological Survey Record 2000/05, Mineral Resources Tasmania.

Doyle, R.B. and Habraken, F.M. (1993) The Distribution of Sodic Soils in Tasmania. *Australian Journal of Soil Research*, 1991, **31**, pp931-47.

Eyre, B.D. and Pepperel, P. (1999) A spatially intensive approach to water quality monitoring in the Rous River catchment, NSW, Australia. *Journal of Environmental Management*, **56**, pp97-108.

Grayson, R.B., Grippel, C.J., Finlayson, B.L. and Hart, B.T. (1997) Catchment-wide impacts on water quality: the use of 'snapshot' sampling during stable flow. *Journal of Hydrology*, **199**, pp121-134.

Grose, C.J. (2003) *Land Degradation and Salinity Risk Investigation in the Coal River Valley, South East Tasmania*. Department of Primary Industries, Water and Environment, Tasmania, Australia.

Hamlet, A.G. (ed) (2002) *Soil Management – A Guide for Tasmanian Farmers*, First Edition, Department of Primary Industries, Water and Environment, Tasmania, Australia.

Harris, G.P. (2001a) Biogeochemistry of nitrogen and phosphorous in Australian catchments, rivers and estuaries: effects of land use and flow regulation and comparisons with global patterns. *Marine and Freshwater Research*, 2001, **52**, pp139-49.

Harris, G. (2001b) *A Nutrient Dynamics Model for Australian Waterways: Land Use, Catchment Biogeochemistry and Water Quality in Australian Rivers, Lakes and Estuaries*, Australia State of the Environment Second Technical Paper Series (Inland Waters), Department of the Environment and Heritage, Canberra.

Hart, B.T., Bailey, P., Edwards, R., Hortle, K., James, K., McMahon, A., Meredith, C. and Swadling, K. (1990) Effects of salinity on river, stream and wetland ecosystems in Victoria, Australia. *Water Research*, Vol **24**, No. 9, pp1103-1117, 1990.

Hart, B.T., Bailey, P., Edwards, R., Hortle, K., James, K., McMahon, A., Meredith, C. and Swadling, K. (1991) A review of the salt sensitivity of the Australian freshwater biota. *Hydrobiologica*, **210**, pp105-144, 1991.

Hatch, D., Goulding, K. and Murphy, D. (2002) Nitrogen. In *Agriculture, Hydrology and Water Quality*, Haygarth, P.M. and Jarvis, S.C. (eds), pp7-27, CABI Publishing, UK.

Hepper-deGryse Team (1991) *Environmental Impact Assessment: Buckland Military Training Area, Volume 2*. Hepper-deGryse Team, Hobart.

Hilderbrand, R.H., Lemly, A.D., Dolloff, C.A. and Harpster, K.L. (1997). Effects of large woody debris placement on stream channels and benthic macroinvertebrates. *Canadian Journal of Aquatic Sciences*, **54**, 931-939.

Horrigan, N., Choy, S., Marshall, J. and Recknagel, F. (2005) Response of stream macroinvertebrates to changes in salinity and the development of a salinity index. *Marine and Freshwater Research*, 2005, **56**, pp825-833.

Jackson, J.E. (2002) Draft Tasmanian Galaxiidae Recovery Plan 2003-2007. Inland Fisheries Service. 78pp.

Jarvie, H.P., Neal, C., Smart, R., Owen, R., Fraser, D., Forbes, I. And Wade, A. (2001) Use of continuous water quality records for hydrograph separation and to assess short-term variability and extremes in acidity and dissolved carbon dioxide for the River Dee, Scotland. *The Science of the Total Environment*, **265** (2001), pp85-98.

Jerie, K., Houshold I. and Peters D. (2003). *Tasmania's river geomorphology: stream character and regional analysis*. Department of Primary Industries, Water and Environment: Hobart, Tasmania.

Kim, J.P., Reid, M.R., Cunningham, R.G. and Hunter, K.A. (1996) Aqueous Chemistry of Major Ions and Trace Metals in the Clutha river, New Zealand. *Marine and Freshwater Research*, **47**, pp919-28.

Kladviko, E.J., Van Scoyoc, G.E., Monke, E.J., Oates, K.M. and Pask, W. (1991) Pesticide and nutrient movement into subsurface tile drains on a silt loam soil in Indiana. *Journal of Environmental Quality*, **20**, pp264-270.

Koehn, J.D. & O' Connor, W.G.O. (1990) *Biological information for management of native freshwater fish in Victoria*. Department of Conservation and Environment, Melbourne.

Koehnken L. (2005) An approach for incorporating geomorphology into environmental flow regimes. In Warfe D.M. (2005). *A Framework for Determining Holistic Environmental Water Requirements in Tasmania*. Draft Technical Report. Department of Primary Industries, Water and Environment: Hobart, Tasmania.

Krasnicki, T., Pinto, R.D. and Read, M.G. (2000). Australia Wide Assessment of River Health; Tasmania Program Final Report. Department of Primary Industries, Water and Environment, Hobart, Technical Report No. WRA 01/2001.

Lake, P.S. (2000) Disturbance, patchiness, and diversity in streams, *Journal of the North American Benthological Society* 19 (4), pp. 573-592.

Lake, P.S. (2003) Ecological effects of perturbation by drought in flowing waters *Freshwater Biology* 48 (7), pp. 1161-1172.

Liston, P and Maher, W. (1997) *Water quality for maintenance of aquatic ecosystems: Appropriate indicators and analysis*, Australia: state of the environment technical Paper Series (Inland Waters), Department of the Environment, Sport and Territories, Canberra.

Lynch, S. (2002) *Modelled Land Capability Classes of Tasmania, Little Swanport 1:100000 map*. Department of Primary Industries, Water and Environment, Tasmania.

Marsh N.A., Stewardson M.J. and Kennard M.J. (2003) *River Analysis Package*. Cooperative Research Centre for Catchment Hydrology, Monash University: Melbourne, Victoria.

McDowall, R.M., (1996). *Freshwater Fishes of South-Eastern Australia*. Reed Books, Hong Kong 247pp.

- Metzling, L. (1993) Benthic macroinvertebrate structure in streams of different salinities. *Australian Journal of Marine and Freshwater Research*, Vol. **44**: 335-351.
- Meyer, S (2002) *Water quality in two small forested catchments in the Warra LTER site, Tasmania: Source of colour and nutrient enrichment*. Masters Thesis, Master of Environmental Management, University of Tasmania.
- Minshall, G. W. (1984) Aquatic Insect Substrate Relationships. *Ecology of Aquatic Insects*. Edited by V. H. Resh and A. M. Rosenberg. Praeger. pp358-400.
- Murray-Darling Basin Ministerial Council (1987) Murray-Darling Basin Environmental Resources Study. Murray Darling Basin Ministerial Council, Canberra.
- Murray-Darling Basin Ministerial Council (1999) *The Salinity Audit of the Murray-Darling Basin*. The Commonwealth, NSW, Vic, SA, QLD, ACT.
- Neal, C. (2001) The water quality of eastern UK rivers: the study of a highly heterogeneous environment. In Land-Ocean Interaction Measuring and modelling fluxes from river basins to the coastal seas. Huntley, D.A., *et al.* (eds) pp69-104 IWA publishing, UK.
- Nielson, D.L., Brock, M.A., Rees, G.N. and Baldwin, D.S. (2003) Effects of increasing salinity on freshwater ecosystems in Australia. *Australian Journal of Botany*, 2003, **51**, pp655-665.
- Parsons, M. and Norris, R. H. (1996) The effect of habitat specific sampling on biological assessment of water quality using a predictive model. *Freshwater Biology*, **36**: 419-434
- Poff, N.L., Allan, J.D., Bain, M.B., Karr, J.R., Prestegard, K.L., Richter, B.D., Sparks, R.E. and Stromberg, J.C. (1997) The natural flow regime: a paradigm for river conservation and restoration. *BioScience* **47**: 769-784.
- Rosenberg, D.M. and Resh, V.H ( eds.) 1993. Freshwater biomonitoring and benthic macroinvertebrates. Chapman and Hall, New York. 488 p.

Rutherford, J.C., Blackett, S., Saito, L. and Davies-Colley, R.J. (1997) Predicting the effects of shade on water temperature in small stream. *New Zealand Journal of Marine and Freshwater Research*, 1997, **31**, pp7-7-721.

Rutherford, J.C., Marsh, A.M., Davies, P.M. and Bunn, S.E. (2004) Effects of patchy shade on stream water temperature: how quickly do small streams heat and cool? *Marine and Freshwater Research*, 2004, **55**, pp737-748.

Salvia, M., Iffly, J.F., Vander Bought, P., Sary, M. and Hoffman, L. (1999) Application of the 'snapshot' methodology to a basin-wide analysis of phosphorous and nitrogen at stable low flow. *Hydrobiologica*, **410**, pp97-102.

Schofield, N.J. & Davies, P.E. 1996, Measuring the health of our rivers, *Water*, May-June 1996, pp39-43.

Sheldon, F., Thoms, M.C. (2006) Relationships between flow variability and macroinvertebrate assemblage composition: Data from four Australian dryland rivers *River Research and Applications* 22 (2), pp. 219-238.

Simpson, J.C. and Norris, R.H. (2000) Biological assessment of river quality: development of AUSRIVAS models and outputs. In: *Assessing the Biological Quality of Freshwaters: RIVPACS and Other Techniques* (Eds J.F. Wright, D.W. Sutcliffe & M.T.Furse), pp. 125-142. Freshwater Biological Association, Ambleside, U.K.

SKM (2004) *Little Swanport River Catchment: Water Balance Model and Scenario Assessment*. Melbourne, Victoria.

UNESCO (1992) *Water Quality Assessment: A Guide To The Use of Biota, Sediments and Water in Environmental Monitoring*. 1st Edition. Chapman and Hall, London.

Wayland, K.G., Long, D.T., Hyndman, D.W., Pijanowski, B.C., Woodhams, S.M. and Sheridan, K.H., (2003) Identifying Relationships between Baseflow Geochemistry and Land Use with Synoptic Sampling and R-Mode Factor Analysis. *Journal of Environmental Quality*, **32**, pp180-190.



Webb A.A. and Erskine W.D. (2003b). Distribution, recruitment, and geomorphic significance of large woody debris in an alluvial forest stream: Tonghi Creek, southeastern Australia. *Geomorphology* **51**: 109-126.

Webb, B.W., Clack, P.D. and Walling, D.E. (2003) Water-air temperature relationships in a Devon river system and the role of flow. *Hydrological Processes*, **17**, pp3069-3084.

Wells, K.F., Gowland, P.N. and Spate, A.P. (1977) *Natural Resources of the Buckland Training Area, Tasmania*. Technical Memorandum 77/14, CSIRO, Division of Land Use Research, Canberra.

Williams K.J. and Potts B.M. (1996). The natural distribution of *Eucalyptus* species in Tasmania. *Tasforests* **8**: 39-165.

Wintle B.C. (2002). *The Ecology of the Riparian Vegetation in Two East Coast River Catchments, Tasmania*. Honours Thesis, School of Geography and Environmental Studies, University of Tasmania. Hobart, Tasmania.

Wright, G.G., Morrice, J.G. and Pugh, K. (1991) North East Scotland River Catchment Nitrate Loading in Relation to Agricultural Intensity, *Chemistry and Ecology*, **5**, pp266-281.

Wright, J. F., Moss, D., Armitage P. D. and Furse, M. T. (1984). A preliminary classification of running-water sites in Great Britain based on macro-invertebrate species and the prediction of community type using environmental data. *Freshwater Biology* **14**: 221-256.