

## DBA *Eucalyptus barberi* forest and woodland

### Community Description:

*Eucalyptus barberi* forest and woodland has a localised distribution in the Eastern Tiers on the central East Coast. It occurs most commonly as mallee-form woodland in small patches on rock plates and rocky ridges within a matrix of *Eucalyptus pulchella* forest and woodland (DPU). The tree height is less than 15 m and usually much lower due to the poor nature of the sites that this community occupies. The solid crown canopy cover is less than 20% in nearly all cases. The understorey is variable, depending on the location, but is often grassy or sedgy; *Lomandra longifolia* is common. Small shrubs also occur; these are commonly from the families Fabaceae and Ericaceae.

### Benchmarks:

Component	Height (m)	Cover (%)	DBH (cm)	Diameter (cm)	Count (#/ha)	Total Length (m/0.1 ha)
<b>Tree Canopy</b>	4	10	-	-	-	-
<b>Large Trees</b>	-	-	20	-	50	-
<b>Logs</b>	-	-	-	≥ 10	-	10
<b>Large Logs</b>	-	-	-	≥ 10	-	-
<b>Organic Litter</b>	-	80	-	-	-	-
<b>Recruitment</b>	Continuous					

Understorey Life Forms	LF code	# Spp	Cover (%)
Immature Canopy Tree	ICT	1	5
Tree (sub-canopy)/Large Shrub	T	1	10
Medium Shrub/Small Shrub	S	5	10
Prostrate and Mat Forming Shrubs (incl. cushion plants)	PS	1	1
Herbs and Orchids	H	3	1
Grasses	G	2	5
Large Sedge/Rush/Sagg/Lily	LSR	1	10
Mosses and Lichens	ML	1	5
<b>Total</b>	<b>8</b>	<b>15</b>	

### Species lists:

Tree Canopy Species	Common Name	Notes
<i>Eucalyptus barberi</i>	barbers gum	
<i>Eucalyptus pulchella</i>	white peppermint	
<i>Callitris rhomboidea</i>	oyster bay pine	

DBA *Eucalyptus barberi* forest and woodland

Typical Understorey Species *	Common Name	LF Code
<i>Bursaria spinosa</i>	prickly box	T
<i>Callistemon pallidus</i>	yellow bottlebrush	T
<i>Leptospermum grandiflorum</i>	autumn teatree	T
<i>Leptospermum scoparium</i>	common teatree	T
<i>Pomaderris elliptica</i>	yellow dogwood	T
<i>Correa reflexa</i>	common correa	S
<i>Epacris impressa</i>	common heath	S
<i>Epacris tasmanica</i>	eastern heath	S
<i>Euryomyrtus ramosissima</i> subsp. <i>ramosissima</i>	rosy heathmyrtle	S
<i>Hakea megadenia</i>	autumn needlebush	S
<i>Hibbertia riparia</i>	erect guineaflower	S
<i>Leptecophylla divaricata</i>	spreading pinkberry	S
<i>Platylobium formosum</i>	handsome flatpea	S
<i>Pultenaea daphnoides</i> var. <i>obcordata</i>	heartleaf bushpea	S
<i>Spyridium obovatum</i>	dustymiller	S
<i>Stenanthemum pimeleoides</i>	propeller plant	S
<i>Astroloma humifusum</i>	native cranberry	PS
<i>Lasiopetalum micranthum</i>	tasmanian velvetbush	PS
<i>Acaena novae-zelandiae</i>	common buzzy	H
<i>Brachyscome aculeata</i>	daisy	H
<i>Leptorhynchus squamatus</i>	scaly buttons	H
<i>Linum marginale</i>	native flax	H
<i>Veronica formosa</i>	speedwell bush	H
<i>Viola</i> spp.	violet	H
<i>Austrodanthonia</i> spp.	wallabygrass	G
<i>Austrostipa</i> spp.	speargrass	G
<i>Poa labillardierei</i>	silver tussockgrass	G
<i>Poa rodwayi</i>	velvet tussockgrass	G
<i>Baloskion australe</i>	southern cordrush	LSR
<i>Gahnia grandis</i>	cutting grass	LSR
<i>Gahnia rodwayi</i>	dwarf sawsedge	LSR
<i>Lepidosperma filiforme</i>	common rapiersedge	LSR
<i>Lomandra longifolia</i>	sagg	LSR
<i>Cheilanthes austrotenuifolia</i>	green rockfern	GF
<i>Pteridium esculentum</i>	bracken	GF

\*This list is provided as a guide only. The species listed are typical of this plant community type but may not necessarily be present.