

DGW *Eucalyptus gunnii* woodland

Community Description:

Eucalyptus gunnii woodland is dominated by *E. gunnii* (both subspecies *gunnii* and subspecies *divaricata*) occasionally growing to 20 m but more commonly 5 –10 m in height. Other less common eucalypt species in this community are *E. coccifera* and *E. delegatensis*. On the most suitable sites, near Great Lake, *E. gunnii* can exceed 20 m in height, as tall straight trees. Broad, branching trees may develop where they are well-spaced. Trees 5 - 10 m high with juvenile foliage retained for many years are characteristic in areas of poor drainage and frost hollows. This is the appropriate benchmark to use in assessing the condition of the listed *Eucalyptus gunnii* woodland community (Schedule 3A, Nature Conservation Act 2002).

Benchmarks:

Component	Height (m)	Cover (%)	DBH (cm)	Diameter (cm)	Count (#/ha)	Total Length (m/0.1 ha)
Tree Canopy	10	40	-	-	-	-
Large Trees	-	-	25	-	50	-
Logs	-	-	-	≥ 10	-	30
Large Logs	-	-	-	≥ 12.5	-	-
Organic Litter	-	10	-	-	-	-
Recruitment	Continuous					

Understorey Life Forms	LF code	# Spp	Cover (%)
Tree (sub-canopy)/Large Shrub	T	3	20
Medium Shrub/Small Shrub	S	6	50
Prostrate and Mat Forming Shrubs (incl. cushion plants)	PS	1	1
Herbs and Orchids	H	5	5
Grasses	G	1	2
Large Sedge/Rush/Sagg/Lily	LSR	3	10
Ground Ferns and fern allies	GF	1	10
Mosses and Lichens	ML	1	5
Total	8	21	

Species lists:

Tree Canopy Species	Common Name	Notes
<i>Eucalyptus gunnii</i>	cider gum	
<i>Eucalyptus coccifera</i>	snow peppermint	
<i>Eucalyptus delegatensis</i>	gumtopped stringybark	
<i>Eucalyptus subcrenulata</i>	alpine yellow gum	

Typical Understorey Species *	Common Name	LF Code
<i>Monotoca submutica</i>	mountain broomheath	T
<i>Oxylobium ellipticum</i>	golden shaggypea	T
<i>Pittosporum bicolor</i>	cheesewood	T
<i>Richea scoparia</i>	scoparia	T
<i>Tasmania lanceolata</i>	mountain pepper	T
<i>Bauera rubioides</i>	wiry bauera	S
<i>Callistemon viridiflorus</i>	prickly bottlebrush	S
<i>Coprosma nitida</i>	mountain currant	S
<i>Cyathodes</i> spp.	cheeseberry	S
<i>Epacris gunnii</i>	coral heath	S
<i>Epacris lanuginosa</i>	swamp heath	S
<i>Grevillea australis</i>	grevillea	S
<i>Hovea montana</i>	mountain purplepea	S
<i>Leptecophylla juniperina</i> subsp. <i>parvifolia</i>	mountain pinkberry	S
<i>Leucopogon montanus</i>	snow beardheath	S
<i>Lomatia polymorpha</i>	mountain guitarplant	S
<i>Monotoca linifolia</i> subsp. <i>algida</i>	alpine nodding broomheath	S
<i>Olearia erubescens</i>	moth daisybush	S
<i>Olearia phlogopappa</i>	dusty daisybush	S
<i>Olearia pinifolia</i>	prickly daisybush	S
<i>Orites revoluta</i>	revolute orites	S
<i>Richea procera</i>	lax candleheath	S
<i>Pultenaea fasciculata</i>	alpine bushpea	PS
<i>Acaena montana</i>	mountain buzzy	H
<i>Acaena novae-zelandiae</i>	common buzzy	H
<i>Asperula gunnii</i>	mountain woodruff	H
<i>Brachyscome spathulata</i>	blue daisy	H
<i>Geranium</i> spp.	geranium	H
<i>Oreomyrrhis ciliata</i>	fringed caraway	H
<i>Rubus gunnianus</i>	alpine raspberry	H
<i>Senecio leptocarpus</i>	western groundsel	H
<i>Stylidium graminifolium</i>	narrowleaf triggerplant	H
<i>Veronica</i> spp.	speedwell	H
<i>Austrodanthonia</i> spp.	wallabygrass	G
<i>Poa gunnii</i>	gunns snowgrass	G
<i>Baloskion australe</i>	southern cordrush	LSR
<i>Gymnoschoenus sphaerocephalus</i>	buttongrass	LSR
<i>Lepidosperma filiforme</i>	common rapiersedge	LSR
<i>Gleichenia microphylla</i>	scrambling coralfern	GF

*This list is provided as a guide only. The species listed are typical of this plant community type but may not necessarily be present.