

DMW Midlands Woodland complex

Community Description:

Midlands woodland complex is a woodland community, only including vegetation with less than 20% solid crown cover. The community comprises the woodland facies of *Eucalyptus ovata* forest and woodland (DOV), *E. viminalis* grassy forest and woodland (DVG) and *E. pauciflora* forest and woodland on dolerite (DPD) where these occur at altitudes of less than 300 m on dolerite and basalt, mainly in the Midlands region of Tasmania. This is the appropriate benchmark to use in assessing the condition of the listed *Eucalyptus ovata* forest and woodland community (Schedule 3A, *Nature Conservation Act 2002*) where it occurs under the above conditions in the Midlands region.

Benchmarks:

Component	Height (m)	Cover (%)	DBH (cm)	Diameter (cm)	Count (#/ha)	Total Length (m/0.1 ha)
Tree Canopy	20	10	-	-	-	-
Large Trees	-	-	80	-	5	-
Logs	-	-	-	≥ 10	-	20
Large Logs	-	-	-	≥ 40	-	-
Organic Litter	-	80	-	-	-	-
Recruitment	Continuous					

Understorey Life Forms	LF code	# Spp	Cover (%)
Immature Canopy Tree	ICT	1	5
Tree (sub-canopy)/Large Shrub	T	2	5
Medium Shrub/Small Shrub	S	4	5
Prostrate and Mat Forming Shrubs (incl. cushion plants)	PS	4	5
Herbs and Orchids	H	8	5
Grasses	G	8	50
Large Sedge/Rush/Sagg/Lily	LSR	1	5
Medium to Small Sedge/Rush/Sagg/Lily	MSR	6	5
Mosses and Lichens	ML	1	5
Total	9	35	

Species lists:

Tree Canopy Species	Common Name	Notes
<i>Eucalyptus ovata</i>	black gum	
<i>Eucalyptus pauciflora</i>	cabbage gum	
<i>Eucalyptus viminalis</i>	white gum	

Last reviewed – 4 July 2022

[Tasmanian Vegetation Monitoring and Mapping Program](#)

Department of Natural Resources and Environment Tasmania

<https://nre.tas.gov.au/>

Typical Understorey Species *	Common Name	LF Code
<i>Acacia dealbata</i>	silver wattle	T
<i>Acacia mearnsii</i>	black wattle	T
<i>Banksia marginata</i>	silver banksia	T
<i>Bursaria spinosa</i>	prickly box	T
<i>Hibbertia</i> spp.	guineaflower	S
<i>Leucopogon virgatus</i>	twiggy beardheath	S
<i>Lissanthe strigosa</i>	peachberry heath	S
<i>Pimelea humilis</i>	dwarf riceflower	S
<i>Acrotriche serrulata</i>	ants delight	PS
<i>Astroloma humifusum</i>	native cranberry	PS
<i>Bossiaea prostrata</i>	creeping bossia	PS
<i>Hibbertia prostrata</i>	prostrate guineaflower	PS
<i>Acaena echinata</i>	spiny sheepsburr	H
<i>Arthropodium milleflorum</i>	pale vanilla-lily	H
<i>Convolvulus angustissimus</i>	blushing bindweed	H
<i>Euchiton collinus</i>	common cottonleaf	H
<i>Gonocarpus tetragynus</i>	common raspwort	H
<i>Hypoxis glabella</i>	tiny yellowstar	H
<i>Lagenophora stipitata</i>	blue bottledaisy	H
<i>Leptorhynchos squamatus</i>	scaly buttons	H
<i>Oxalis perennans</i>	grassland woodsorrel	H
<i>Plantago</i> spp.	plantain	H
<i>Solenogyne</i> spp.	flat-herb	H
<i>Viola hederacea</i>	ivyleaf violet	H
<i>Wahlenbergia</i> spp.	bluebell	H
<i>Agrostis</i> spp.	bent	G
<i>Austrodanthonia</i> spp.	wallabygrass	G
<i>Austrostipa</i> spp.	speargrass	G
<i>Deyeuxia</i> spp.	bentgrass	G
<i>Dichelachne</i> spp.	plumegrass	G
<i>Ehrharta stipoides</i>	weeping grass	G
<i>Elymus scaber</i>	rough wheatgrass	G
<i>Poa</i> spp.	tussockgrass	G
<i>Themeda triandra</i>	kangaroo grass	G
<i>Dianella revoluta</i>	spreading flaxlily	LSR
<i>Lomandra longifolia</i>	sagg	LSR
<i>Carex breviculmis</i>	shortstem sedge	MSR
<i>Centrolepis</i> spp.	bristlewort	MSR
<i>Lepidosperma</i> spp.	swordsedge	MSR
<i>Schoenus apogon</i>	common bogsedge	MSR

*This list is provided as a guide only. The species listed are typical of this plant community type but may not necessarily be present.