

DRI *Eucalyptus risdonii* forest and woodland

Community Description:

Eucalyptus risdonii forest and woodlands are dominated by a sparse canopy of *E. risdonii* trees often with a mallee habit. They are dry sclerophyll communities with sparse medium and low shrub layers and an often-grassy ground layer. This is the appropriate benchmark to use in assessing the condition of the listed *Eucalyptus risdonii* forest and woodland community (Schedule 3A, Nature Conservation Act 2002).

Benchmarks:

Component	Height (m)	Cover (%)	DBH (cm)	Diameter (cm)	Count (#/ha)	Total Length (m/0.1 ha)
Tree Canopy	3	50	-	-	-	-
Large Trees	-	-	10	-	200	-
Logs	-	-	-	≥ 10	-	10
Large Logs	-	-	-	≥ 10	-	-
Organic Litter	-	20	-	-	-	-
Recruitment	Continuous					

Understorey Life Forms	LF code	# Spp	Cover (%)
Immature Canopy Tree	ICT	1	10
Tree (sub-canopy)/Large Shrub	T	4	20
Medium Shrub/Small Shrub	S	6	15
Prostrate and Mat Forming Shrubs (incl. cushion plants)	PS	4	15
Herbs and Orchids	H	2	1
Grasses	G	5	20
Large Sedge/Rush/Sagg/Lily	LSR	3	10
Mosses and Lichens	ML	1	1
Total	8	26	

Species lists:

Tree Canopy Species	Common Name	Notes
<i>Eucalyptus risdonii</i>	risdon peppermint	
<i>Eucalyptus amygdalina</i>	black peppermint	
<i>Eucalyptus tenuiramis</i>	silver peppermint	
<i>Eucalyptus viminalis</i>	white gum	

Typical Understorey Species *	Common Name	LF Code
<i>Acacia dealbata</i>	silver wattle	T
<i>Acacia mearnsii</i>	black wattle	T
<i>Acacia verticillata</i>	prickly moses	T
<i>Allocasuarina littoralis</i>	black sheoak	T
<i>Allocasuarina verticillata</i>	drooping sheoak	T
<i>Bursaria spinosa</i>	prickly box	T
<i>Cassinia aculeata</i>	dollybush	T
<i>Dodonaea viscosa</i>	broadleaf hopbush	T
<i>Exocarpos cupressiformis</i>	native cherry	T
<i>Pomaderris elliptica</i>	yellow dogwood	T
<i>Acacia genistifolia</i>	spreading wattle	S
<i>Acacia myrtifolia</i>	redstem wattle	S
<i>Correa reflexa</i>	common correa	S
<i>Daviesia latifolia</i>	hop bitterpea	S
<i>Daviesia ulicifolia</i>	spiky bitterpea	S
<i>Epacris impressa</i>	common heath	S
<i>Lissanthe strigosa</i>	peachberry heath	S
<i>Ozothamnus obcordatus</i>	yellow everlastingbush	S
<i>Philotheca virgata</i>	twiggy waxflower	S
<i>Pultenaea daphnoides</i> var. <i>obcordata</i>	heartleaf bushpea	S
<i>Pultenaea gunnii</i>	golden bushpea	S
<i>Pultenaea juniperina</i>	prickly beauty	S
<i>Tetratheca labillardierei</i>	glandular pinkbells	S
<i>Acrotriche serrulata</i>	ants delight	PS
<i>Astroloma humifusum</i>	native cranberry	PS
<i>Bossiaea prostrata</i>	creeping bossia	PS
<i>Pultenaea pedunculata</i>	matted bushpea	PS
<i>Gonocarpus tetragynus</i>	common raspwort	H
<i>Opercularia varia</i>	variable stinkweed	H
<i>Stylidium graminifolium</i>	narrowleaf triggerplant	H
<i>Wahlenbergia</i> spp.	bluebell	H
<i>Agrostis</i> spp.	bent	G
<i>Austrodanthonia</i> spp.	wallabygrass	G
<i>Austrostipa</i> spp.	speargrass	G
<i>Deyeuxia</i> spp.	bentgrass	G
<i>Elymus scaber</i>	rough wheatgrass	G
<i>Poa rodwayi</i>	velvet tussockgrass	G
<i>Dianella revoluta</i>	spreading flaxlily	LSR
<i>Diplarrena moraea</i>	white flag-iris	LSR
<i>Gahnia radula</i>	thatch sawsedge	LSR
<i>Lepidosperma laterale</i>	variable sword-sedge	LSR
<i>Lomandra longifolia</i>	sagg	LSR

*This list is provided as a guide only. The species listed are typical of this plant community type but may not necessarily be present.