

DTD *Eucalyptus tenuiramis* forest and woodland on dolerite: (forest)

Community Description:

Eucalyptus tenuiramis forest and woodland on dolerite is dominated by *E. tenuiramis* trees rarely more than 25 m in height, and considerably smaller on highly insolated sites. These dry sclerophyll communities are generally characterised by shrubby understoreys with high cover and diversity and a sparse ground layer. This benchmark is one of 2 benchmarks available to assess the condition of DTD.

Benchmarks:

Component	Height (m)	Cover (%)	DBH (cm)	Diameter (cm)	Count (#/ha)	Total Length (m/0.1 ha)
Tree Canopy	25	30	-	-	-	-
Large Trees	-	-	60	-	15	-
Logs	-	-	-	≥ 10	-	40
Large Logs	-	-	-	≥ 30	-	-
Organic Litter	-	80	-	-	-	-
Recruitment	Continuous					

Understorey Life Forms	LF code	# Spp	Cover (%)
Immature Canopy Tree	ICT	1	5
Tree (sub-canopy)/Large Shrub	T	1	5
Medium Shrub/Small Shrub	S	14	35
Prostrate and Mat Forming Shrubs (incl. cushion plants)	PS	1	1
Herbs and Orchids	H	3	1
Grasses	G	2	2
Large Sedge/Rush/Sagg/Lily	LSR	4	15
Ground Ferns and fern allies	GF	1	1
Scramblers/Climbers/Epiphytes	SCE	1	1
Mosses and Lichens	ML	1	5
Total	10	29	

Species lists:

Tree Canopy Species	Common Name	Notes
<i>Eucalyptus tenuiramis</i>	silver peppermint	
<i>Eucalyptus amygdalina</i>	black peppermint	
<i>Eucalyptus obliqua</i>	stringybark	
<i>Eucalyptus pulchella</i>	white peppermint	

DTD *Eucalyptus tenuiramis* forest and woodland on dolerite: (forest)

Typical Understorey Species *	Common Name	LF Code
<i>Acacia dealbata</i>	silver wattle	T
<i>Allocasuarina</i> spp.	sheoak	T
<i>Banksia marginata</i>	silver banksia	T
<i>Bedfordia salicina</i>	tasmanian blanketleaf	T
<i>Callitris rhomboidea</i>	oyster bay pine	T
<i>Exocarpos cupressiformis</i>	common native-cherry	T
<i>Hakea lissosperma</i>	mountain needlebush	T
<i>Leptospermum grandiflorum</i>	autumn teatree	T
<i>Leptospermum scoparium</i>	common teatree	T
<i>Melaleuca squarrosa</i>	scented paperbark	T
<i>Acacia</i> spp.	wattle	S
<i>Correa reflexa</i>	common correa	S
<i>Epacris</i> spp.	heath	S
<i>Hibbertia riparia</i>	erect guineaflower	S
<i>Leptomeria drupacea</i>	erect currantbush	S
<i>Leucopogon collinus</i>	white beardheath	S
<i>Lissanthe strigosa</i>	peachberry heath	S
<i>Lomatia tinctoria</i>	guitarplant	S
<i>Platylobium triangulare</i>	arrow flatpea	S
<i>Pultenaea</i> spp.	bushpea	S
<i>Spyridium</i> spp.	dustymiller	S
<i>Tetratheca pilosa</i>	hairy pinkbells	S
<i>Xanthorrhoea</i> spp.	grasstree	S
<i>Acrotriche serrulata</i>	ants delight	PS
<i>Astroloma humifusum</i>	native cranberry	PS
<i>Goodenia lanata</i>	trailing native-primrose	H
<i>Gonocarpus teucrioides</i>	forest raspwort	H
<i>Helichrysum scorpioides</i>	curling everlasting	H
<i>Austrodanthonia</i> spp.	wallabygrass	G
<i>Austrostipa</i> spp.	speargrass	G
<i>Ehrharta distichophylla</i>	hairy ricegrass	G
<i>Dianella</i> spp.	flaxlily	LSR
<i>Diplarrena moraea</i>	white flag-iris	LSR
<i>Gahnia</i> spp.	sawsedge	LSR
<i>Gymnoschoenus sphaerocephalus</i>	buttongrass	LSR
<i>Lepidosperma</i> spp.	swordsedge	LSR
<i>Lomandra longifolia</i>	sagg	LSR
<i>Pteridium esculentum</i>	bracken	GF
<i>Billardiera longiflora</i>	purple appleberry	SCE
<i>Cassytha pubescens</i>	downy dodderlaurel	SCE
<i>Comesperma volubile</i>	blue lovecreeper	SCE

*This list is provided as a guide only. The species listed are typical of this plant community type but may not necessarily be present.