

## Wet Eucalyptus Forest and Woodland

### WOB *Eucalyptus obliqua* forest with broadleaf shrubs

#### Community Description:

*Eucalyptus obliqua* forest with broadleaf shrubs are tall to very tall forest communities typically with an understorey composed of broadleaved shrubs, the most common including *Pomaderris apetala*, *Nematolepis squamea* and *Olearia argophylla*, with a high proportion of ground ferns. This community is widely distributed in moderate to high rainfall areas on a variety of substrates.

#### Benchmarks:

Component	Cover %	Height (m)	DBH (cm)	#/ha	Length (m)/0.1 ha
Canopy	30%	-	-	-	-
Large Trees	-	40	90	18	-
Organic Litter	90%	-	-	-	-
Logs	-	-	≥ 10	-	40
Large Logs	-	-	≥ 45	-	-
Recruitment	Episodic				

Understorey Life Forms	LF code	# Spp	Cover %
Tree or large shrub	T	7	40
Shrub	S	6	5
Herbs and orchids	H	4	1
Grass	G	2	1
Large sedge/rush/sagg/lily	LSR	2	5
Ground fern	GF	3	10
Tree fern	TF	1	5
Scrambler/Climber/Epiphytes	SCE	3	1
Mosses and Lichens	ML	1	5
<b>Total</b>	<b>9</b>	<b>29</b>	

#### Species lists:

Canopy Tree Species	Common Name	Notes
<i>Eucalyptus obliqua</i>	stringybark	
<i>Eucalyptus amygdalina</i>	black peppermint	
<i>Eucalyptus brookeriana</i>	brookers gum	
<i>Eucalyptus dalrympleana</i>	mountain white gum	
<i>Eucalyptus delegatensis</i>	gumtopped stringybark	
<i>Eucalyptus globulus</i>	Tasmanian blue gum	
<i>Eucalyptus nitida</i>	western peppermint	
<i>Eucalyptus regnans</i>	giant ash	
<i>Eucalyptus viminalis</i>	white gum	

WOB *Eucalyptus obliqua* forest with broadleaf shrubs

Typical Understorey Species *	Common Name	LF Code
<i>Acacia</i> spp.	wattle	T
<i>Anopterus glandulosus</i>	tasmanian laurel	T
<i>Banksia marginata</i>	silver banksia	T
<i>Bedfordia salicina</i>	tasmanian blanketleaf	T
<i>Cenarrhenes nitida</i>	native plum	T
<i>Monotoca glauca</i>	goldey wood	T
<i>Nematolepis squamea</i>	satinwood	T
<i>Olearia</i> spp.	daisybush	T
<i>Phyllocladus aspleniifolius</i>	celerytop pine	T
<i>Pomaderris</i> spp.	dogwood	T
<i>Prostanthera lasianthos</i>	christmas mintbush	T
<i>Zieria arborescens</i>	stinkwood	T
<i>Bauera rubioides</i>	wiry bauera	S
<i>Coprosma</i> spp.	currant	S
<i>Leucopogon</i> spp.	beardheath	S
<i>Olearia viscosa</i>	viscid daisybush	S
<i>Oxylobium ellipticum</i>	golden shaggypea	S
<i>Pimelea drupacea</i>	cherry riceflower	S
<i>Pultenaea</i> spp.	bushpea	S
<i>Acaena novae-zelandiae</i>	common buzzy	H
<i>Dryophila cyanocarpa</i>	turquoise berry	H
<i>Geranium potentilloides</i>	mountain cranesbill	H
<i>Hydrocotyle hirta</i>	hairy pennywort	H
<i>Chiloglottis</i> spp.	orchids	H
<i>Ranunculus</i> spp.	buttercup	H
<i>Senecio</i> spp.	groundsel	H
<i>Viola hederacea</i>	ivyleaf violet	H
<i>Deyeuxia</i> spp.	bentgrass	G
<i>Ehrharta</i> spp.	ricegrass	G
<i>Festuca</i> spp.	fescue	G
<i>Dianella tasmanica</i>	forest flaxlily	LSR
<i>Gahnia grandis</i>	cutting grass	LSR
<i>Lepidosperma elatius</i>	tall swordsedg	LSR
<i>Blechnum</i> spp.	waterfern	GF
<i>Histiopteris incisa</i>	batswing fern	GF
<i>Hypolepis</i> spp.	groundfern	GF
<i>Polystichum proliferum</i>	mother shieldfern	GF
<i>Dicksonia antarctica</i>	soft treefern	TF
<i>Billardiera longiflora</i>	purple appleberry	SCE
<i>Clematis</i> spp.	clematis	SCE
<i>Microsorium pustulatum</i>	kangaroo fern	SCE

\*This list is provided as a guide only. The species listed are typical of this plant community type but may not necessarily be present.